

BRIUMVI (UBLITUXIMAB-XIY) ORDER FORM

PATIENT INFORMATION

Patient Name: _____ DOB: _____

Mobile Number: _____ Patient Weight: _____

Allergies: _____

DIAGNOSIS (Provider must specify)

MS: G35

Other: _____

PROVIDER INFORMATION

Provider Name (print name): _____ Provider NPI: _____

Signature: _____ Date: _____

Contact Name: _____ Phone: _____ Fax: _____

Email Address: _____

Prerequisites to treatment – ensure the following information is complete and attached with referral:

- Demographics Labs and tests supporting diagnosis Office/progress notes

PRE-MEDICATION (Not typically indicated)

Acetaminophen (Tylenol) 500 mg PO

Famotidine 20 mg IV

Methylprednisolone (Solu-Medrol) 125 mg IVP

Benadryl 25mg PO

Cetirizine (Zyrtec) 10 mg PO

Other: _____ Dose: _____ Route: _____

MEDICATION

MEDICATION	DOSE / ROUTE / FREQUENCY
Briumvi	<input type="checkbox"/> First infusion: 150 mg IV; Second infusion: 450 mg IV two weeks after first infusion, then 450 mg IV 24 weeks after first infusion and every 24 weeks thereafter <input type="checkbox"/> 450 mg IV every 24 weeks

New Start Therapy Continuation of Therapy Date of last dose (if applicable): _____

LABS / SPECIAL INSTRUCTIONS