Local Infusion

MYLOCALINFUSION.COM

PEMGARDA (PEMIVIBART) ORDER FORM

PATIENT INFORMATION

Patient Name: _____ DOB: _____

Mobile Number: _____ Patient Weight: _____

Allergies: _____

DIAGNOSIS (Provider must specify)

❑ Pre-exposure prophylaxis of coronavirus disease 2019 (COVID-19) in adults and adolescents (12 years of age) who have moderate-to-severe immune compromise, ICD 10: _____

PROVIDER INFORMATION

Provider Name (print name): _____ Provider NPI: _____

Signature: _____ Date: _____

Contact Name: _____ Phone: _____ Fax: _____

Email Address: _____

Prerequisites to treatment – ensure the following information is complete and attached with referral:

❑ Demographics ❑ Labs and tests supporting diagnosis ❑ Office/progress notes

PRE-MEDICATION (Not typically indicated)

❑ Acetaminophen (Tylenol) 500 mg PO

❑ Benadryl 25mg PO

❑ Famotidine 20 mg IV

❑ Cetirizine (Zyrtec) 10 mg PO

❑ Methylprednisolone (Solu-Medrol) 125 mg IVP

❑ Other: _____ Dose: _____ Route: _____

MEDICATION

MEDICATION	DOSE	ROUTE	FREQUENCY
Pemgarda	❑ Initial Dose: 4500 mg administered as a single intravenous infusion. ❑ Repeat Dose: 4500 mg administered as a single intravenous infusion approximately every 3 months.	❑ IV	❑ q 3 months

❑ New Start Therapy ❑ Continuation of Therapy Date of last dose (if applicable): _____

LABS / SPECIAL INSTRUCTIONS

FAX NUMBERS: ❒ CT: 203.433.0621 ❒ ME: 207.407.7272 ❒ NH: 603.217.5371 ❒ NJ: 201.581.4521

Order valid for 1 year from date of signature unless otherwise specified here: _____