

PEMGARDA (PEMIVIBART) ORDER FORM

PATIENT INFORMATION

Patient Name: _____ DOB: _____

Mobile Number: _____ Patient Weight: _____

Allergies: _____

DIAGNOSIS (Provider must specify)

Pre-exposure prophylaxis of coronavirus disease 2019 (COVID-19) in adults and adolescents (12 years of age) who have moderate-to-severe immune compromise, ICD 10: _____

PROVIDER INFORMATION

Provider Name (print name): _____ Provider NPI: _____

Signature: _____ Date: _____

Contact Name: _____ Phone: _____ Fax: _____

Email Address: _____

Prerequisites to treatment – ensure the following information is complete and attached with referral:

- Demographics Labs and tests supporting diagnosis Office/progress notes

PRE-MEDICATION (Not typically indicated)

- Acetaminophen (Tylenol) 500 mg PO Famotidine 20 mg IV Methylprednisolone (Solu-Medrol) 125 mg IVP
- Benadryl 25mg PO Cetirizine (Zyrtec) 10 mg PO
- Other: _____ Dose: _____ Route: _____

MEDICATION

MEDICATION	DOSE	ROUTE	FREQUENCY
Pemgarda	<input type="checkbox"/> Initial Dose: 4500 mg administered as a single intravenous infusion. <input type="checkbox"/> Repeat Dose: 4500 mg administered as a single intravenous infusion approximately every 3 months.	<input type="checkbox"/> IV	<input type="checkbox"/> q 3 months

New Start Therapy Continuation of Therapy Date of last dose (if applicable): _____

LABS / SPECIAL INSTRUCTIONS